

Genetic Diversity and Host Range of *Verticillium dahliae* Isolates from Artichoke and Other Vegetable Crops in Spain

M. Berbegal, Instituto Agroforestal Mediterráneo, Universidad Politécnica de Valencia, Camino de Vera s/n, 46022 Valencia, Spain; **A. Ortega**, Departamento de Producción Vegetal y Microbiología, Escuela Politécnica Superior de Orihuela, Universidad Miguel Hernández, Ctra. Beniel km 3.2, 03312 Orihuela, Alicante, Spain; **M. M. Jiménez-Gasco**, Department of Plant Pathology, The Pennsylvania State University, University Park 16802; **C. Olivares-García** and **R. M. Jiménez-Díaz**, Departamento de Agronomía, ETSIAM, Universidad de Córdoba, and Instituto de Agricultura Sostenible, CSIC, Campus Rabanales, Edificio C-4 'Celestino Mutis', 14071 Córdoba, Spain; and **J. Armengol**, Instituto Agroforestal Mediterráneo, Universidad Politécnica de Valencia

ABSTRACT

Berbegal, M., Ortega, A., Jiménez-Gasco, M. M., Olivares-García, C., Jiménez-Díaz, R. M., and Armengol, J. 2010. Genetic diversity and host range of *Verticillium dahliae* isolates from artichoke and other vegetable crops in Spain. *Plant Dis.* 94:396-404.

Artichoke is severely affected by *Verticillium* wilt, caused by *Verticillium dahliae*, in eastern-central Spain, which is one of the most important vegetable-cropping areas in the country. To determine genetic and virulence variability in local populations of *V. dahliae*, 18 isolates collected from artichoke and other vegetable species cultivated in eastern-central Spain were selected to represent local vegetative compatibility groups (VCGs). Diversity in the isolates was characterized by molecular markers and virulence in 12 important hosts for that region. Recently developed microsatellite markers (simple-sequence repeats) and polymorphic sequences were used to assess the genetic variation among those isolates to reveal any association occurring among host source, VCG, and virulence. Although all isolates caused severe disease symptoms on artichoke, cardoon, eggplant, and watermelon, those from artichoke had a limited host range and isolates from watermelon, muskmelon, and eggplant were not pathogenic to some of the hosts tested. VCG diversity was related to differential virulence in certain hosts.

Verticillium wilts are economically important diseases of dicotyledonous crops worldwide and are caused by the soilborne pathogen *Verticillium dahliae* Kleb. (35). *Verticillium* wilt negatively affects artichoke production (*Cynara cardunculus* L. var. *scolymus* (L.) Fiori) in the Comunidad Valenciana region located in eastern-central Spain, one of the most important vegetable-cropping areas in the country (1,4). Knowledge about the genetic and virulence diversity in local populations of *V. dahliae* associated with different crops is a key component for the management of *Verticillium* wilt diseases, particularly through the use of host resistance and choice of crop rotation in a given region (12,36,44,46). Studies during the last decade have shown that genetic diversity in *V. dahliae* populations is higher than previously expected (39). Also, severe disease outbreaks may emerge in a region on vegetable species previously considered to be nonhosts of *V. dahliae* (2,6,7,10,24,36,39).

Studies also have demonstrated that *V. dahliae* from certain hosts shows a degree of host specificity and differential virulence (7,19,38). For instance, *V. dahliae* isolates from cocoa and pepper have limited host ranges (7,38,45), and cotton isolates differing in virulence (defined as the amount of disease caused in a host genotype) to cotton cultivars show differential virulence on other herbaceous crops (28,41).

Genetic diversity within *V. dahliae* populations has been studied mainly by means of vegetative compatibility grouping and molecular analyses (10,20,26). Isolates that are vegetatively compatible belong to the same vegetative compatibility group (VCG) based on their genetically controlled ability to undergo hyphal anastomosis and form stable heterokaryons (23,29). For strictly asexually reproducing fungi, hyphal fusion is necessary for genetic exchange among different isolates (35). Therefore, *V. dahliae* isolates from different VCGs are thought to be genetically isolated populations which may vary in some ecological, physiological, and virulence traits (12,23,39). A correlation between virulence and the VCG of *V. dahliae* isolates has been demonstrated in certain hosts (5,16,22,26,46,48). Conversely, *V. dahliae* isolates in the same VCG but from different hosts may differ in host

range, suggesting that a pathogenicity trait might be determined by selection pressure established by cropping practices (7,12,45,48).

Studies on the relationships between VCGs, virulence, and molecular diversity within *V. dahliae* often yielded contradictory results. Bhat and Subbarao (6) reported that host specificity and virulence diversity were not associated to random amplified polymorphic DNA (RAPD) patterns. In contrast, RAPD markers clearly differentiated between *V. dahliae* isolates from woody and herbaceous hosts (33). Similarly, RAPD markers demonstrated correlation between molecular grouping, VCGs, and virulence among isolates from different hosts (47,48).

The diversity in VCG and VCG-associated virulence among artichoke *V. dahliae* isolates from the main artichoke-growing areas in the Comunidad Valenciana region was recently investigated (20). Four VCGs (VCG1A, VCG2A, VCG2B and VCG4B) were identified associated with affected artichokes. VCG2B was the predominant group. Using a large sample of isolates representative of each VCG, it was concluded that VCG2B and VCG4B isolates were the most virulent to artichoke whereas isolates in VCG1A and a single heterokaryon self-incompatible (HSI) isolate were the least virulent to that host. Furthermore, specific polymerase chain reaction (PCR) markers and amplified fragment length polymorphism (AFLP) analysis demonstrated that a correlation existed between molecular differences among artichoke isolates in VCG2B and virulence to artichoke (10,20). Also, isolates of VCG2B were subdivided into two subgroups (VCG2Ba and VCG2Br) based on complementation to reference isolates (20). VCG2Br isolates were defined by the positive complementation of nitrate-nonutilizing (*nit*) mutants of tested isolates with reference strains from the international Ohio Agricultural Research and Development Center (OARDC) and Israel (26,39). Conversely, VCG2Ba isolates were characterized by positive complementation of *nit1* mutants of tested isolates with *nit* mutants of VCG2B isolates from

Corresponding author: M. Berbegal
E-mail: mobermar@etsia.upv.es

Accepted for publication 30 October 2009.

doi:10.1094/PDIS-94-4-0396
© 2010 The American Phytopathological Society

artichoke but failed to establish positive complementation with VCG2B reference strains from the international OARDC and Israel (20).

The objectives of this study were to (i) comparatively determine the host range of *V. dahliae* isolates from artichoke and other vegetable crops commonly grown in eastern-central Spain and (ii) determine the genetic variation among those isolates that may reveal any association occurring among host source, VCG, and virulence.

MATERIALS AND METHODS

***V. dahliae* isolates.** Eighteen monocolonial *V. dahliae* isolates collected from vegetable production areas of eastern-central Spain were used in this study (Table 1). VCGs of artichoke isolates were determined in an earlier study by Jiménez-Díaz et al. (20). VCG typing of non-artichoke isolates was performed by complementation tests of *nit* mutants using international reference isolates as previously described (25). Artichoke isolates were selected to represent the VCG diversity identified in the Comunidad Valenciana region (20). Non-artichoke isolates were from vegetable crops representative of a wide diversity of commonly grown host species that can be used in rotations with artichoke in the area of the study (Table 1). Isolates were stored in a 15% glycerol solution at -80°C in 1.5-ml cryovials in the culture collection of the Grupo de Investigación en Hongos Fito-patógenos, Instituto Agroforestal Mediterráneo, Valencia.

DNA extraction and specific PCR assay. DNA was extracted from mycelia of 7-day-old cultures of *V. dahliae* isolates grown in potato-dextrose broth (Sigma-Aldrich, St. Louis) incubated at room temperature on an orbital shaker. Approximately 20 mg of freeze-dried mycelium was ground to a fine powder with a sterile mortar and pestle. DNA was extracted from ground mycelia using the E.Z.N.A. Plant DNA Kit (Omega Bio-Tek, USA). DNA concentration was determined by comparison of different dilutions of samples with the molecular marker Lambda DNA (New England Biolabs, Ipswich, MA). *V. dahliae* isolates were characterized by duplex- or multiplex-PCR assays using several primer pairs as previously described (10,30). Primer pair DB19/DB22 (9) yields *V. dahliae*-specific polymorphic DNA bands of 539 or 523 bp (30). The primer pairs INTD2f/INTD2r and INTND2f/INTND2r produced PCR markers of 462 and 824 bp, previously associated with the cotton- and olive-defoliating (D) and -nondefoliating (ND) *V. dahliae* pathotypes, respectively (30–32). In contrast, primer pair DB19/espdef01 amplifies a 334-bp marker present in D isolates as well as in some ND isolates of VCG1B and VCG2B (10,11).

Amplification of polymorphic markers and sequencing. Four simple-

sequence repeats (SSRs) and seven previously developed polymorphic sequences (3) were used to characterize diversity of *V. dahliae* isolates in this study (Table 2). PCR reaction mixtures (25 μl) contained 2.5 μl of 10 \times standard buffer (15 mM of MgCl_2) (Gene Choice, Frederick, MD) 0.25 μM each primer (Table 2) (Nucleic Acid Facility, Penn State University), 250 μM each dNTP, 1.25 U of *Taq* DNA polymerase (Gene Choice), and 10 to 50 ng of template DNA. The initial denaturation step (94°C , 4 min) was followed by 40 cycles of denaturation at 94°C for 1 min, annealing at T_a (Table 2) for 1 min, and elongation at 72°C for 1 min, with a final elongation step at 72°C for 6 min. Amplifications were performed on a Mastercycler[®] ep (Eppendorf AG, Hamburg, Germany). Microsatellite alleles were separated on 6% polyacrylamide gels in 0.5 \times Tris-borate-EDTA (TBE) buffer and visualized with UV light after ethidium bromide staining. A 25-bp ladder (Invitrogen, Carlsbad, CA) was used to estimate molecular weight of PCR products. Allele sizes and variation of repeats were also assessed by sequence analysis of amplified SSR loci and polymorphic sequences. Purified amplicons were sequenced in both directions using the primers, and sequences of alleles at each locus were aligned and compared using Sequencher v. 4.7 software (Gene Codes, Ann Arbor, MI).

Combined datasets of SSR and polymorphic sequences were developed by

assigning simple number codes to each allele per locus for all isolates. Cluster analysis was done using simple matching coefficient by the unweighted pair-group method using arithmetic averages (UP-GMA). Calculations were performed using SIMQUAL and SAHN programs of the software package NTSYS-pc (version 2.02j; Exeter Software, Setauket, NY).

Pathogenicity tests. Two experiments (I and II) were performed to test the 18 selected *V. dahliae* isolates (Table 1) for virulence on artichoke cv. Madrigal and the following cultivars of 11 vegetable crop species commonly grown in eastern-central Spain in rotation with artichoke: cardoon (*Cynara cardunculus* var. *cardunculus* cv. Blanco Lleno Mejorado), cauliflower (*Brassica oleracea* cv. Bola de Nieve X), cucumber (*Cucumis sativus* cv. Negrito), eggplant (*Solanum melongena* cv. Black Beauty), lettuce (*Lactuca sativa* cv. Romana Larga Rubia), muskmelon (*Cucumis melo* cv. Sancho), pepper (*Cap-sicum annuum* cv. Dulce de España), potato (*S. tuberosum* cv. Spunta), pumpkin (*Cucurbita pepo* cv. Dulce de Horno), tomato (*S. lycopersicum* cv. Marmandino), and watermelon (*Citrullus lanatus* cv. Dulce Maravilla). Certified seed of these Verticillium wilt nonresistant cultivars were obtained from commercial seed companies. Experiment I included 12 *V. dahliae* isolates from artichoke representative of the VCG diversity found among artichoke isolates in the Comunidad Valenciana.

Table 1. Isolates of *Verticillium dahliae* used in this study, listed by host source, geographic origin, vegetative compatibility group (VCG), and molecular marker

Isolate	Host	Origin (province)	VCG ^b	Molecular marker (bp) ^a		
				334 (D)	824 (ND)	462 (D)
V403	Artichoke	Valencia	1A	+	–	+
V404	Artichoke	Valencia	2A	–	+	–
V542	Artichoke	Alicante	2A	–	+	–
V536	Artichoke	Castellón	2Br	–	+	–
V549	Artichoke	Valencia	2Br	+	–	–
V593	Artichoke	Alicante	2Br	–	+	–
V613	Artichoke	Castellón	2Br	+	–	–
V574	Artichoke	Alicante	2Ba	+	–	–
V675	Artichoke	Valencia	2Ba	+	–	–
V684	Artichoke	Valencia	4B	–	+	–
V687	Artichoke	Valencia	4B	–	+	–
V547	Artichoke	Alicante	HSI	+	–	–
V2	Watermelon	Valencia	2A	–	+	–
V10	Pumpkin	Valencia	2A	–	+	–
V20	Muskmelon	Castellón	2A	–	+	–
V25	Tomato	Valencia	2A	–	+	–
V61	Potato	Alicante	4B	–	+	–
V158	Eggplant	Valencia	4B	–	+	–

^a Amplicons produced in polymerase chain reaction assays using primer sets DB19/DB22/espdef01 (334 bp), INTD2f/INTD2r (462 bp), and INTND2f/INTND2r (824 bp). Amplicons of 334 and 462 bp were previously associated with the cotton-defoliating pathotype, and an amplicon of 824 bp with the cotton-nondefoliating pathotype (10,30). D = defoliating, ND = nondefoliating, + = positive amplification, and – = negative amplification.

^b VCG2Br is defined by positive complementation of nitrate-nonutilizing (*nit*) mutants of the tested isolates with the complementary mutants of international Ohio Agricultural Research and Development Center reference and Israeli tester strains (25,39). VCG2Ba is defined by positive complementation of *nit1* mutants of the tested isolates with selected NitM mutants of isolates from artichoke that previously had been assigned to VCG2Br (20). Isolates assigned to VCG2Ba failed to establish positive complementation with *nit* mutants of the international reference and Israeli tester strains. HSI = heterokaryon self incompatible.

ana region (20). In experiment II, six *V. dahliae* isolates from different vegetable species grown in the area of study were tested and the artichoke V593 isolate representative of the predominant VCG, VCG2B, was included for comparison (Table 1).

In the two experiments, plants were inoculated by the root-dip inoculation method. Surface-disinfested (1% NaClO, 2 min), germinated seed were sown in trays filled with a sterile (autoclaved twice, 120°C for 30 min) potting mixture (peat/sand, 2:1, vol/vol). Plants were

grown in a greenhouse at 20 to 25°C until inoculation (two to three true leaves). Inocula were obtained from 3- to 4-week-old cultures on potato-dextrose agar (PDA) (Biokar Diagnostics, Beauvais, France) by flooding them with sterile distilled water and scraping their surface with a sterile

Table 2. Polymorphic molecular markers used to characterize *Verticillium dahliae* isolates

Locus ^a	Primer sequence (5'-3')	T _a (°C) ^b	Size range (bp)	SNP location and type ^c	Na ^d
VdSSR1	F: TTGATCTAGGAAAGCGCATGAC R: CCAGGCAACAATGCAGACA	55	120–128	—	2
VdSSR2	F: CCCCAGCCTCTCTTCTTCTAT R: AGCAGAATCACTCATAGGGTGC	47	245–260	—	4
VdSSR4	F: TCAGAACCGAAGTAACCAA R: GAATCACGCTCCTTCACGGTA	47	222–224	—	2
VdSSR5	F: GCAGGGGGTGGCCTCAGCACT R: CGCAGGCATGCCTGGGTGTG	59	217–224	—	2
VdSNP1	F: TGACACTGAAAAGGATACACCG R: TGGAAACACCCATAAGACAACG	55	—	78 C/G; 98 C/T; 181 C/T; 189 T/C; 222 G/C; 292 A/T;	4
VdSNP2	F: GATCATGAAGCGAGTGCAGAAGAAC R: TTGGGGCGTGTGGAAGGAT	59	—	328 C/T; 329 A/G; 334 G/C; 348 C/G; 386 T/G	8
VdSNP3	F: TGCAGCGAAGACAGAGAC R: CTAAGCCACCCAGCAACACA	59	—	80 C/T; 313 T/C	3
VdSNP4	F: GAAGGCGTCCACCCAGTCGTC R: GCTAATCGCCCCGTCGCAACA	50	—	288 A/G; 289 —/G	3
VdSNP5	F: ACGGGGGATGGGTGTGAT R: TTCTGGCAAGTTTGACATGACTG	59	—	223 T/A	2
VdSNP6	F: GCGGCGGCATGCATCACCTC R: CCATGGCCTCGGGCGTTCG	59	—	204 A/T	2
VdSNP7	F: TCAGCTGCGCAACTGCTCAA R: TCACCGCTTTCCGCTACTACCTT	55	—	82 C/—; 109 T/C; 113 A/T	3

^a VSSR codes, microsatellite markers; VSNP codes, polymorphic sequences.

^b T_a, annealing temperature used in polymerase chain reaction amplifications.

^c Nucleotide position and changes in the GenBank accessioned sequence; —, gap in the nucleotide sequence.

^d Number of alleles observed.




Fig. 1. Dendrogram of 18 *Verticillium dahliae* isolates from artichoke and other vegetable crops based on cluster analysis of the simple matching similarity coefficient by unweighted pair-group method using arithmetic averages (UPGMA) of the sequence variations detected by four microsatellite markers and seven polymorphic sequences. Artichoke isolates: V403, V574, V549, V675, V613, V547, V404, V542, V684, V687, V536, and V593. Watermelon isolate: V2. Pumpkin isolate: V10. Muskmelon isolate: V20. Tomato isolate: V25. Potato isolate: V61. Eggplant isolate: V158.

spatula. Inoculum suspensions were filtered through sterile cheesecloth and the conidial concentration in the filtrate was adjusted to 10^6 conidia ml^{-1} for each isolate. Plants were uprooted from the trays and their roots washed thoroughly under tap water to remove substrate residues, slightly trimmed, and dipped for 10 min in the conidia suspension of an isolate. Non-inoculated control plants were treated the same as the inoculated ones but were dipped in sterile distilled water. Inoculated and control plants were transplanted into 12-cm-diameter plastic pots filled with the sterile potting mixture (one plant per pot) and incubated at the same conditions described above in the greenhouse for 1 month. There were 10 replicated plants for each isolate-cultivar combination in the experiments.

Disease reactions were assessed 1 month after inoculation by the severity of symptoms in individual plants scored on a 1-to-6 scale according to percentage of affected tissue (1 = no symptoms; 2 = 1 to 25, 3 = 26 to 50, 4 = 51 to 75, and 5 = 76 to 100% of tissue affected; and 6 = dead plant). Symptoms included chlorosis, defoliation, stunting, and wilting. At the end of the experiments, isolations were made from symptomatic and nonsymptomatic plants to confirm infection and colonization by the pathogen. Plant shoots were surface disinfested (1% NaClO, 2 min) and small internal fragments of vascular tissue were plated onto PDA supplemented with streptomycin sulfate (PDAS) at 0.5 mg ml^{-1} . Cultures were incubated

and *V. dahliae* colonies identified as described previously (4).

Statistical analysis. Experiments I and II consisted of a factorial treatment design (*V. dahliae* isolates \times host cultivars) with 10 replicates, each one consisting of a single potted plant, in a complete randomized design. The experiments were repeated once. A two-tailed *F* test indicated that there was no heterogeneity of variances between the two repetitions of each experiment, allowing data for each treatment to be combined for statistical analyses. Combined datasets were analyzed using a nonparametric procedure for the analysis of ordinal data in factorial experiments (8,42). The overall effect of *Verticillium* isolates and VCGs on disease reaction on each inoculated host cultivar was determined by the analysis of variance type statistic of ranked data. The PROC mixed procedure in SAS (version 9.0; SAS Institute, Cary, NC) was used to generate relative effects (REs), and the LD_CI macro was used to generate 95% confidence intervals (8,42). RE values are generated by the equation: $\text{RE} = (R - 0.5)/N$; where *R* is the mean treatment ranking and *N* is the total number of observations in the analysis. Linear single-degree-of-freedom contrasts were computed to test the effect of selected experimental treatment combinations (18).

RESULTS

Vegetative compatibility of *V. dahliae* isolates. Of the six non-artichoke *V. dahliae* isolates characterized to VCG in this

study, isolates from watermelon (V2), pumpkin (V10), muskmelon (V20), and tomato (V25) were assigned to VCG2A. Isolates from potato (V61) and eggplant (V158) were assigned to VCG4B. Isolate V61 formed weak complementation with the VCG4B reference isolate.

Molecular characterization of *V. dahliae* isolates by specific PCR, SSR, and polymorphic sequences analyses. All 18 *V. dahliae* isolates used in this study amplified the species-specific 539- or 523-bp polymorphic DNA bands in PCR assays using the DB19/DB22 primer pair. PCR assays of isolates using primer pairs DB19/espdef01 (334-bp amplicon), INTD2f/INTD2r (462-bp amplicon), and INTND2f/INTND2r (824-bp amplicon) produced one of the following patterns of PCR products: 334 (+), 824 (–), 462 (+); 334 (–), 824 (+), 462 (–); and 334 (+), 824 (–), 462 (–) (Table 1). The only artichoke isolate representing VCG1A amplified the 334- and the 462-bp DNA fragments. Isolates assigned to VCG2A and VCG4B amplified the 824-bp marker regardless of host source. Artichoke isolates of VCG2Br amplified either the 824-bp marker (designated as VCG2Br₈₂₄) or the 334-bp marker (designated as VCG2Br₃₃₄). However, artichoke isolates of VCG2Ba amplified the 334-bp marker (designated as VCG2Ba₃₃₄) as well as the HSI isolate from artichoke.

Based on the four SSR and seven polymorphic sequences in the 18 isolates, there was a total of 35 alleles (Table 2). Loci produced between two and eight alleles.

Table 3. Median (M) and mean rankings (R) calculated for the severity of *Verticillium* wilt symptoms on artichoke, cardoon, eggplant, muskmelon, potato, tomato, and watermelon caused by isolates of *Verticillium dahliae* from artichoke of different vegetative compatibility groups (VCGs)^a

VCG, isolates ^b	Artichoke		Cardoon		Eggplant		Muskmelon		Potato		Tomato		Watermelon	
	M	R	M	R	M	R	M	R	M	R	M	R	M	R
Control	1.0	5.5	1.0	12.6	1.0	9.5	1.0	33.0	1.0	25.5	1.0	43.5	1.0	10.5
1A, V403	5.4	81.0	4.1	95.4	1.8	25.4	1.2	42.4	1.0	25.5	1.1	49.7	4.9	90.3
2A, V404 and V542	4.3	49.9	2.3	48.8	3.0	51.3	2.0	67.6	2.0	64.3	2.4	90.8	4.0	66.6
2Ba ₃₃₄ , V574 and V675	4.8	59.6	2.5	54.4	2.6	43.3	2.8	91.7	2.1	67.8	1.3	60.3	4.6	76.2
2Br ₃₃₄ , V549 and V613	4.3	48.8	2.4	51.2	4.0	77.5	3.0	92.0	2.3	79.7	1.6	75.9	3.8	54.9
2Br ₈₂₄ , V536 and V593	5.9	101.6	4.4	100.0	5.0	109.5	1.4	48.9	2.3	76.2	1.2	53.1	4.8	87.9
4B, V684 and V687	4.7	56.8	2.4	49.5	4.0	81.3	1.6	53.3	1.9	63.9	1.1	49.8	2.8	45.6
HSI, V547	3.9	32.7	3.0	65.4	4.4	90.5	2.0	69.0	1.6	49.8	1.0	43.5	4.3	67.8
Contrast (<i>P</i>) ^c														
1A vs all	0.0013		<0.0001		<0.0001		0.0001		<0.0001		0.0614		<0.0001	
2A vs all	0.0618		0.0020		<0.0001		0.8730		0.6699		<0.0001		0.6327	
2Ba ₃₃₄ vs all	0.6052		0.0088		<0.0001		<0.0001		0.3647		0.9785		0.2265	
2Br ₃₃₄ vs all	0.0162		0.0025		0.0549		0.0002		0.0025		0.0216		0.0133	
2Br ₈₂₄ vs all	<0.0001		<0.0001		<0.0001		0.0024		0.0080		0.1399		<0.0001	
4B vs all	0.3326		0.0415		0.0588		0.0604		0.6869		0.0145		0.0039	
HSI vs all	<0.0001		0.9227		0.0046		0.7775		0.2325		<0.0001		0.7992	
2A vs 4B	0.4195		0.9508		0.0005		0.1900		0.9688		<0.0001		0.0790	
HSI vs 2Br ₃₃₄ , 2Ba ₃₃₄	0.0012		0.2763		0.0020		0.0494		0.0439		<0.0001		0.8266	
2Ba ₃₃₄ vs 2Br ₃₃₄ , 2Br ₈₂₄	0.0011		0.0002		<0.0001		0.0081		0.2681		0.5732		0.4621	
2Ba ₃₃₄ vs 2Br ₃₃₄	0.1102		0.6293		<0.0001		0.9688		0.2479		0.1025		0.0112	
2Br ₈₂₄ vs 2Br ₃₃₄ , 2Ba ₃₃₄	<0.0001		<0.0001		<0.0001		<0.0001		0.7483		0.0365		0.0001	

^a For median disease rating (M), severity of *Verticillium* wilt symptoms was assessed visually according to the percentage of foliar tissue affected (foliar symptoms including chlorosis, defoliation, stunting, and wilting) using an ordinal 1-to-6 rating scale in which 1 = no symptoms; 2 = 1 to 25, 3 = 26 to 50, 4 = 51 to 75, and 5 = 76 to 100% of foliar tissue affected; and 6 = dead plant.

^b VCG of the isolates inoculated. Isolates were inoculated on the different species by root immersion; roots of control plants were immersed in sterile distilled water. HSI = heterokaryon self incompatible.

^c Linear single-degree-of-freedom contrast computed to test the effect of selected treatment combinations. Probability for the *t* statistic of linear single-degree-of-freedom contrasts, significance level *P* < 0.05.

UPGMA analysis of the SSR and polymorphic sequence data sets separated the isolates into six clusters (designated A, B, C, D, E, and F) based on similarity coefficient values of approximately 0.30 to 0.75, which correlated with VCGs of isolates (Fig. 1). Cluster A comprised the only isolate representative of VCG1A. Cluster B included five VCG2B isolates from artichoke that amplified the 334-bp marker and the single HSI isolate. Cluster C included isolates in VCG2B that amplified the 824-bp marker. Cluster D contained all VCG2A isolates from artichoke and other hosts and one potato isolate in VCG4B (V61). Cluster E contained isolates in VCG4B from artichoke that shared a similarity coefficient of 0.77 to 1. Finally, the eggplant isolate V158 in VCG4B was separated from the rest in cluster F.

Pathogenicity of *V. dahliae* isolates.

All 18 *V. dahliae* isolates were pathogenic on artichoke, cardoon, eggplant, and watermelon as indicated by mean symptom severity rankings significantly greater than those in the noninoculated controls ($P < 0.05$) but differed in virulence on specific hosts. Conversely, only some of the isolates in the study were pathogenic to muskmelon, potato, and tomato (Tables 3 and 4; Figs. 2 and 3). No symptoms developed on inoculated cauliflower, cucumber, lettuce, or pepper plants (*data not shown*).

Therefore, all 18 isolates were considered nonpathogenic to these species. *V. dahliae* was isolated from symptomatic plants of susceptible species and only occasionally recovered from nonsymptomatic plants of cauliflower, cucumber, pepper, and pumpkin.

Median and mean rankings for the severity of symptoms caused by artichoke isolates on different hosts in experiment I, as well as results of linear single-degree-of-freedom contrasts computed to test the effect of selected treatment combinations, are shown in Table 3. Estimated relative effects and confidence intervals are represented in Figure 2. Significant differences in mean symptom severity rankings were observed among hosts ($P < 0.0001$), between VCGs of isolates ($P < 0.0001$), and for the VCG–host interaction ($P < 0.0001$) in experiment I.

Collectively, artichoke *V. dahliae* isolates of VCG1A and VCG2Br₈₂₄ were significantly more virulent to artichoke, cardoon, and watermelon than those in other VCGs, as indicated by mean symptom severity rankings (Table 3). Isolates of VCG2Br₈₂₄ and the HSI isolate were significantly more virulent on eggplant than those in the other VCGs (Table 3).

Among artichoke isolates of VCG2B, those in VCG2Br₈₂₄, as a group, were significantly more virulent on artichoke, car-

doon, and eggplant ($P < 0.0001$) than isolates in VCG2Br₃₃₄ and VCG2Ba₃₃₄ (Table 3; Fig. 2). Conversely, isolates in VCG2Ba₃₃₄ and VCG2Br₃₃₄ were significantly more virulent on muskmelon than those in the other VCGs (Table 3). However, artichoke isolates in VCG1A (V403), VCG2Br₈₂₄ (V536 and V593), VCG4B (V684 and V687), and the HSI isolate (V547) were not pathogenic to muskmelon (Table 3; Fig. 2). Isolates of VCG2Br₃₃₄ ($P = 0.0025$) and VCG2Br₈₂₄ ($P = 0.0080$) were significantly more virulent on potato than those in the other VCGs but isolates in VCG1A was not pathogenic (Table 3). Isolates of VCG2A and VCG2Br₃₃₄ were significantly more virulent on tomato than those in other VCGs, which were considered not pathogenic to tomato (Table 3; Fig. 2). VCG2B isolates that amplified the 334-bp marker (VCG2Br₃₃₄ and VCG2Ba₃₃₄) and the HSI isolate were compared for virulence because they grouped together in cluster B (Fig. 1). Significant differences were detected among those VCG subgroups in virulence to all hosts except for cardoon, muskmelon, and watermelon (Table 3).

Median and mean rankings for the severity of symptoms caused by isolates on different hosts in experiment II, as well as results of linear single-degree-of-freedom contrast, are shown in Table 4. Estimated

Table 4. Median (M) and mean rankings (R) calculated for the severity of Verticillium wilt symptoms on artichoke, cardoon, eggplant, muskmelon, potato, tomato and watermelon caused by isolates of *Verticillium dahliae* collected from different vegetable crops^a

Group, isolates ^b	Artichoke		Cardoon		Eggplant		Muskmelon		Potato		Tomato		Watermelon	
	M	R	M	R	M	R	M	R	M	R	M	R	M	R
Control	1.0	5.0	1.0	6.0	1.0	5.5	1.0	28.0	1.0	13.0	1.0	27.0	1.0	9.5
Cucurbitaceous														
V2	4.9	42.9	3.9	48.1	3.3	28.5	2.0	48.9	2.1	44.5	1.0	27.0	3.0	21.6
V10	4.9	41.6	3.8	47.5	4.2	46.7	1.5	41.0	1.9	38.2	2.2	58.0	6.0	47.0
V20	2.8	15.6	1.9	19.5	2.4	17.3	1.7	44.1	1.2	18.9	1.0	27.0	4.3	33.3
Solanaceous														
V25	5.0	44.2	3.8	45.8	3.9	40.1	1.3	38.5	2.7	59.2	1.8	44.8	5.3	42.3
V61	5.7	61.2	4.7	58.5	5.3	68.9	1.3	38.5	2.2	47.5	2.4	64.6	5.3	41.3
V158	5.2	49.4	2.8	33.6	4.3	48.5	1.3	39.7	2.4	50.4	1.0	27.0	5.7	41.1
Artichoke														
V593	5.5	56.6	3.7	46.9	5.0	65.0	1.2	32.9	1.9	39.2	1.3	35.6	4.7	35.9
Contrast (P) ^c														
V2 vs all	0.6695		0.2871		<0.0001		0.1959		0.6627		<0.0001		0.0017	
V10 vs all	0.4464		0.4214		0.5854		0.9302		0.4966		0.0006		<0.0001	
V20 vs all	<0.0001		<0.0001		<0.0001		0.5540		<0.0001		<0.0001		0.4181	
V25 vs all	0.9464		0.6172		0.1414		0.6906		0.0001		0.5038		0.2701	
V593 vs all	0.0148		0.3881		<0.0001		0.1102		0.3271		0.3358		0.7383	
V61 vs all	<0.0001		<0.0001		<0.0001		0.6906		0.3566		<0.0001		0.3039	
V158 vs all	0.3519		0.0151		0.2752		0.8752		0.1549		<0.0001		0.3199	
VCG2A vs VCG4B ^d	<0.0001		<0.0001		<0.0001		0.4593		<0.0001		<0.0001		0.2444	
Cucurbitaceous vs solanaceous ^e	<0.0001		0.0645		<0.0001		0.2561		0.0002		0.0476		0.0582	
Cucurbitaceous vs artichoke	0.0017		0.1676		<0.0001		0.0750		0.2542		0.7821		0.7474	
Solanaceous vs artichoke	0.4255		0.8794		<0.0001		0.3197		0.0081		0.1454		0.3523	

^a For median disease rating (M), severity of Verticillium wilt symptoms was assessed visually according to the percentage of foliar tissue affected (foliar symptoms including chlorosis, defoliation, stunting, and wilting) using an ordinal 1-to-6 rating scale in which 1 = no symptoms; 2 = 1 to 25, 3 = 26 to 50, 4 = 51 to 75, and 5 = 76 to 100% of foliar tissue affected; and 6 = dead plant.

^b Isolates inoculated grouped by the taxonomic relationship of the isolates' host source. Cucurbitaceous = isolates from watermelon (V2), pumpkin (V10) and muskmelon (V20). Solanaceous = isolates collected from tomato (V25), potato (V61), and eggplant (V158). Artichoke = isolate from artichoke (V593). Isolates were inoculated on the different species by root immersion; roots of control plants were immersed in sterile distilled water.

^c Linear single-degree-of-freedom contrast computed to test the effect of selected treatment combinations. Probability for the t statistic of linear single-degree-of-freedom contrasts, significance level $P < 0.05$.

^d Linear single-degree-of-freedom contrast of the mean rankings corresponding to isolates in VCG2A (V2, V10, V20, and V25) and VCG4B (V61).

^e Linear single-degree-of-freedom contrast of the mean rankings corresponding to groups of isolates based on the taxonomic relationship of the isolates' host source.

relative effects and confidence intervals are represented in Figure 3. Significant differences in mean symptom severity rankings were observed among hosts ($P < 0.0001$), among isolates differing in host source (P

< 0.0001), and for the isolate–host interaction ($P < 0.0001$) in experiment II.

The artichoke isolate V593 was significantly ($P = 0.0148$) more virulent on artichoke and eggplant compared with all

others isolates (Table 4). Similarly, the pumpkin isolate V10 was the most virulent isolate on watermelon (Table 4) but it failed to cause disease on pumpkin (*data not shown*). None of the isolates were




Fig. 2. Estimated relative effects (REs) in experiment I based on the analysis of variance-type statistics of ranked data for the severity of *Verticillium* wilt symptoms on **A**, artichoke; **B**, cardoon; **C**, muskmelon; **D**, watermelon; **E**, potato; **F**, tomato; and **G**, eggplant caused by *Verticillium dahliae* artichoke isolates of vegetative compatibility groups (VCGs) 1A, 2A, 2Ba₃₃₄, 2Br₃₃₄, 2Br₈₂₄, 4B, and heterokaryon self-incompatible (HSI). Confidence intervals (95%) for relative effects are represented by vertical lines.

pathogenic on muskmelon (Table 4; Fig. 3), and muskmelon isolate V20 was not pathogenic to potato (Table 4; Fig. 3). The potato isolate V61 was significantly more virulent on artichoke, cardoon, eggplant,

and tomato compared with all other isolates (Table 4). Only isolates from pumpkin (V10), tomato (V25), and potato (V61) were pathogenic to tomato (Table 4; Fig. 3).

Isolates in VCG2A (V2, V10, V20, and V25) and VCG4B (V61), which grouped together in cluster C, differed in virulence on all hosts except muskmelon and watermelon (Table 4). A degree of association




Fig. 3. Estimated relative effects (RE) in experiment II based on the analysis of variance-type statistics of ranked data for the severity of *Verticillium* wilt symptoms on **A**, artichoke; **B**, cardoon; **C**, muskmelon; **D**, watermelon; **E**, potato; **F**, tomato; and **G**, eggplant caused by *Verticillium dahliae* isolates: V2 = watermelon isolate, V10 = pumpkin isolate, V20 = muskmelon isolate, V25 = tomato isolate, V593 = artichoke isolate, V61 = potato isolate, and V158 = eggplant isolate. Confidence intervals (95%) for relative marginal effects are represented by vertical lines. Bars are textured according to the taxonomic relationship of the isolates' host source: isolates collected from cucurbitaceous plants (gray-shaded bar), solanaceous plants (white bar), and artichoke (cross-hatched bar).

was found between the taxonomic relationship of host source of isolates and virulence to given hosts. Thus, isolates from solanaceous plants (V25, V61, and V158) were significantly more virulent on potato than all other isolates (Table 4; Fig. 3), isolates from cucurbitaceous plants (V2, V10, and V20) were significantly less virulent on artichoke compared with other isolates, and the artichoke isolate V593 was significantly more virulent on eggplant than other isolates (Table 4; Fig. 3).

DISCUSSION

Genetic and pathogenic characterization of *V. dahliae* isolates in an agricultural region may provide important insights into the epidemiology and management of the diseases they cause. In this study, *V. dahliae* isolates collected from artichoke and other vegetable species cultivated in eastern-central Spain were characterized by means of molecular markers and virulence to 12 important hosts for that region. We demonstrate that differential virulence occurs among isolates from certain hosts and that differences can be associated with genetic diversity among isolates. Although all isolates caused severe disease symptoms on artichoke, cardoon, eggplant, and watermelon, those from artichoke showed a limited host range and isolates from watermelon, muskmelon, and eggplant were not pathogenic to some of the hosts tested. A previous report demonstrated that eggplant isolates exhibit host specificity and differential virulence on other hosts, whereas isolates from artichoke, potato, tomato, and watermelon did not (7).

Isolates of *V. dahliae* are usually considered to be host adapted rather than host specific, because isolates seem to be most virulent on certain hosts but often have the potential to infect a wide range of hosts (16,38). In general, isolates of *V. dahliae* are highly virulent on the hosts from which they were originally isolated (7,13,19,38). In our study, all isolates were pathogenic on hosts other than the host source. However, our results showed a possible association between the taxonomic relationship of the host source of isolates and their virulence on certain hosts.

Other authors have suggested that host range is more influenced by crop history than the original host source of the isolate (45) or by the VCG of the isolate (48). Several studies demonstrated that VCGs of *V. dahliae* do not appear to be related to pathogenicity of particular host species but there is some correlation with virulence to certain hosts such as VCG4A and potato in the United States (39). In this study, *V. dahliae* isolates from artichoke differed in virulence on certain hosts, and the virulence appeared related to VCG. Thus, artichoke isolates in VCG2A caused severe disease in tomato, which agreed with earlier reports describing isolates in VCG2A as virulent on tomato (46). In our study,

the artichoke isolate in VCG1A was not pathogenic to muskmelon and potato but highly virulent on artichoke. In contrast, artichoke isolates in VCG1A were less virulent on the host source than on cotton (20). In previous studies, individual isolates within VCGs varied considerably with respect to virulence (17,19,43). However, there was a consistent relationship between VCG and virulence on the host source, which was reported for artichoke (20) and potato and cotton (27,43).

Artichoke has been described as the cultivated form of cardoon (40). To our knowledge, there are no references about the incidence of Verticillium wilt in cultivated cardoon, and no disease has been observed on cardoon crops grown in fields in the Comunidad Valenciana region of Spain, which is highly infested with *V. dahliae* (M. Berbegal and J. Armengol, unpublished). Therefore, we included cardoon in pathogenicity tests because cardoon germplasm may be a potential source of resistance to *V. dahliae* in artichoke breeding programs. However, our results revealed that cardoon is very susceptible to the pathogen, to an extent similar to artichoke and eggplant (14,37,38).

In strictly asexually reproducing pathogens, such as *V. dahliae*, host-associated differences in neutral molecular markers can be investigated with the assumption that, on particular hosts, associations between loci controlling pathogenicity and neutral molecular loci might emerge (34). Combining neutral molecular markers with pathogenicity assays represents a useful method of studying host specificity in pathogens with broad host ranges. Based on specific PCR markers, VCG2B from artichoke in eastern-central Spain is genetically heterogeneous, and that subgrouping of isolates associated with such heterogeneity correlates with virulence to artichoke and cotton (20). In our study, subgrouping among artichoke isolates in VCG2B based on genetic and molecular differences was also correlated with virulence to different hosts. The pattern of correlation was similar on artichoke, cardoon, and eggplant, to which isolates of VCG2Br₈₂₄ were the most virulent. Artichoke isolates of VCG2Br₈₂₄ were also the most virulent to potato and watermelon.

Qin et al. (36) recently reported the existence of genetically distinct subpopulations of *V. dahliae* adapted to specific hosts but each being cross pathogenic to various crops. Some of the past inconsistencies in the associations between *V. dahliae* isolates and molecular subgroupings (5,7) may have resulted from pathogenicity tests with a limited number of hosts, underestimating the full pathogenic potential of the tested isolates (36).

The correlation between neutral genetic markers and VCGs in *V. dahliae* isolates from artichoke and other crops has already been demonstrated using AFLPs (10). In

the present study, recently developed microsatellite (SSRs) and polymorphic sequences were used to easily characterize molecular variability of *V. dahliae* isolates from artichoke and other vegetable species grown in the Comunidad Valenciana region. Results showed that *V. dahliae* isolates within a VCG shared most of the alleles and could be clearly separated from those in other VCGs regardless of their host source. UPGMA analysis clustered isolates in accordance with their VCG and confirmed molecular heterogeneity of artichoke isolates in VCG2B based on AFLPs and specific PCR markers (10,20). Molecular variability among isolates could be associated with VCG diversity and differential virulence on certain hosts.

Occasionally, weak complementation occurs between strains of different VCGs that indicate transitory functional heterokaryosis before incompatibility is expressed, and suggests that VCGs might not be as completely isolated in genetic terms as previously thought. According to some authors, weak heterokaryon reactions often occur between testers or *nit* mutant isolates of VCG2 and VCG4 (21,43). In our study, isolates in VCG2A and 4B clustered very closely in the dendrogram and one of the isolates from potato that formed weak complementation reaction with the VCG4B reference isolate grouped with the VCG2A isolates in the same cluster. These results suggest that isolates in VCG2A and VCG4B could be more similar to each other than to isolates of other VCGs. This agrees with results of Collado-Romero et al., who found a similar relationship by AFLP analysis (10).

According to Bhat and Subbarao, results of pathogenicity tests in the greenhouse are predictive of host ranges in the field (7). Therefore, decisions concerning crop rotations in eastern-central Spain can be based on data reported here (20). Moreover, rotation with nonhost crops might reduce the probability for the selection of highly virulent strains of *V. dahliae* favored by monoculture (15). All isolates studied here were nonpathogenic to cauliflower, cucumber, lettuce, pepper, and pumpkin and, thus, their use in rotation with susceptible crops would not likely increase inoculum in the soil. However, recommendations for cropping sequences should be made with caution, and further research on isolates from a wider variety of vegetable crops is needed.

ACKNOWLEDGMENTS

M. Berbegal was a recipient of an FPU fellowship from the Ministry of Education and Science of Spain (AP2002-3818). We thank A. Vicent for helpful comments on the manuscript prior to submission.

LITERATURE CITED

1. Armengol, J., Berbegal, M., Giménez-Jaime, A., Romero, S., Beltrán, R., Vicent, A., Ortega, A., and García-Jiménez, J. 2005. Incidence of Verticillium wilt of artichoke in eastern Spain

- and role of inoculum sources on crop infection. *Phytoparasitica* 33:397-405.
2. Berbegal, M., and Armengol, J. 2009. First report of *Verticillium* wilt of faba bean caused by *Verticillium dahliae* in Spain. *Plant Dis.* 93:432.
3. Berbegal, M., Garzón, C., Ortega, A., Armengol, J., Jiménez-Díaz, R. M., and Jiménez-Gasco, M. M. Development and application of new molecular markers for analysis of genetic diversity in *Verticillium dahliae* populations. *Plant Pathol.* In press.
4. Berbegal, M., Ortega, A., García-Jiménez, J., and Armengol, J. 2007. Inoculum density-disease development relationship in *Verticillium* wilt of artichoke caused by *Verticillium dahliae*. *Plant Dis.* 91:1131-1136.
5. Bhat, R. G., Smith, R. F., Koike, S. T., Wu, B. M., and Subbarao, K. V. 2003. Characterization of *Verticillium dahliae* isolates and wilt epidemics of pepper. *Plant Dis.* 87:789-797.
6. Bhat, R. G., and Subbarao, K. V. 1999. First report of *Verticillium dahliae* causing artichoke wilt in California. *Plant Dis.* 83:782.
7. Bhat, R. G., and Subbarao, K. V. 1999. Host range specificity in *Verticillium dahliae*. *Phytopathology* 89:1218-1225.
8. Brunner, E., Domhof, S., and Langer, F. 2002. Nonparametric Analysis of Longitudinal Data in Factorial Experiments. John Wiley & Sons, New York.
9. Carder, J. H., Morton, A., Tabrett, A. M., and Barbara, D. J. 1994. Detection and differentiation by PCR of subspecific groups within two *Verticillium* species causing vascular wilts in herbaceous hosts. Pages 91-97 in: *Modern Assays for Plant Pathogenic Fungi*. A. Schots, F. M. Dewey, and R. Oliver, eds. CAB International, Wallingford, UK.
10. Collado-Romero, M., Mercado-Blanco, J., Olivares-García, C., Valverde-Corredor, A., and Jiménez-Díaz, R. M. 2006. Molecular variability within and among *Verticillium dahliae* vegetative compatibility groups determined by fluorescent amplified fragment length polymorphism and polymerase chain reaction markers. *Phytopathology* 96:485-495.
11. Collins, A., Mercado-Blanco, J., Jiménez-Díaz, R. M., Olivares, C., Clewes, E., and Barbara, D. J. 2005. Correlation of molecular markers and biological properties in *Verticillium dahliae* and the possible origins of some isolates. *Plant Pathol.* 54:549-557.
12. Chandelier, A., Laurent, F., Dantinne, D., Mariage, L., Etienne, M., and Cavellier, M. 2003. Genetic and molecular characterization of *Verticillium dahliae* isolates from woody ornamentals in Belgian nurseries. *Eur. J. Plant Pathol.* 109:943-952.
13. Chang, R. J., and Eastburn, D. M. 1994. Host range of *Verticillium dahliae* from horseradish and pathogenicity of strains. *Plant Dis.* 78:503-506.
14. Cherrab, M., Serrhini, M. N., and Charest, P. M. 2000. Characterization of Moroccan isolates of *Verticillium dahliae* Kleb. using RAPD markers. *J. Phytopathol.* 148:243-249.
15. Davis, J. R., Pavek, J. J., Corsini, D. L., Sorensen, L. H., Schneider, A. T., Everson, D. O., Westermann, D. T., and Huisman, O. C. 1994. Influence of continuous cropping of several potato clones on the epidemiology of *Verticillium* wilt of potato. *Phytopathology* 84:207-214.
16. Douhan, L. I., and Johnson, D. A. 2001. Vegetative compatibility and pathogenicity of *Verticillium dahliae* from spearmint and peppermint. *Plant Dis.* 85:297-302.
17. Elena, K. 1999. Genetic relationships among *Verticillium dahliae* isolates from cotton in Greece based on vegetative compatibility. *Eur. J. Plant Pathol.* 105:609-616.
18. Gómez, K. A., and Gómez, A. A. 1984. Statistical Procedures for Agricultural Research, 2nd ed. John Wiley and Sons, New York.
19. Goud, J. C., and Termorshuizen, A. J. 2002. Pathogenicity and virulence of the two Dutch VCGs of *Verticillium dahliae* to woody ornamentals. *Eur. J. Plant Pathol.* 108:771-782.
20. Jiménez-Díaz, R. M., Mercado-Blanco, J., Olivares-García, C., Collado-Romero, M., Bejarano-Alcázar, J., Rodríguez-Jurado, D., Giménez-Jaime, A., García-Jiménez, J., and Armengol, J. 2006. Genetic and virulence diversity in *Verticillium dahliae* populations infecting artichoke in eastern-central Spain. *Phytopathology* 96:288-298.
21. Joaquim, T. R., and Rowe, R. C. 1990. Reassessment of vegetative compatibility relationships among strains of *Verticillium dahliae* using nitrate-nonutilizing mutants. *Phytopathology* 80:1160-1166.
22. Joaquim, T. R., and Rowe, R. C. 1991. Vegetative compatibility and virulence strains of *Verticillium dahliae* from soil and potato plant. *Phytopathology* 81:552-558.
23. Katan, T. 2000. Vegetative compatibility in populations of *Verticillium*—an overview. Pages 69-86 in: *Advances in Verticillium: Research and Disease Management*. Proc. 7th Int. *Verticillium* Symp. E. C. Tjamos, R. C. Rowe, J. B. Heale, and D. R. Fravel, eds. The American Phytopathological Society, St. Paul, MN.
24. Koike, S. T., Subbarao, K. V., Davis, R. M., Gordon, T. R., and Hubbard, J. C. 1994. *Verticillium* wilt of cauliflower in California. *Plant Dis.* 78:1116-1121.
25. Korolev, N., and Katan, T. 1997. Improved medium for selecting nitrate nonutilizing (*nit*) mutants of *Verticillium dahliae*. *Phytopathology* 87:1067-1070.
26. Korolev, N., Katan, J., and Katan, T. 2000. Vegetative compatibility groups of *Verticillium dahliae* in Israel: their distribution and association with pathogenicity. *Phytopathology* 90:529-566.
27. Korolev, N., Pérez-Artés, E., Bejarano-Alcázar, J., Rodríguez-Jurado, D., Katan, J., Katan, T., and Jiménez-Díaz, R. M. 2001. Comparative study of genetic diversity and pathogenicity among populations of *Verticillium dahliae* from cotton in Spain and Israel. *Eur. J. Plant Pathol.* 107:443-456.
28. Korolev, N., Pérez-Artés, E., Mercado-Blanco, J., Bejarano-Alcázar, J., Rodríguez-Jurado, D., Jiménez-Díaz, R. M., Katan, T., and Katan, J. 2008. Vegetative compatibility of cotton-defoliating *Verticillium dahliae* in Israel and its pathogenicity to various crop plants. *Eur. J. Plant Pathol.* 122:603-617.
29. Leslie, J. F. 1993. Fungal vegetative compatibility. *Annu. Rev. Phytopathol.* 31:127-150.
30. Mercado-Blanco, J., Rodríguez-Jurado, D., Parrilla-Araujo, S., and Jiménez-Díaz, R. M. 2003. Simultaneous detection of the defoliating and nondefoliating *Verticillium dahliae* pathotypes in infected olive plants by duplex, nested polymerase chain reaction. *Plant Dis.* 87:1487-1494.
31. Mercado-Blanco, J., Rodríguez-Jurado, D., Pérez-Artés, E., and Jiménez-Díaz, R. M. 2001. Detection of the nondefoliating pathotype of *Verticillium dahliae* in infected olive plants by nested PCR. *Plant Pathol.* 50:1-12.
32. Mercado-Blanco, J., Rodríguez-Jurado, D., Pérez-Artés, E., and Jiménez-Díaz, R. M. 2002. Detection of the defoliating pathotype of *Verticillium dahliae* in infected olive plants by nested PCR. *Eur. J. Plant Pathol.* 108:1-13.
33. Nigro, F., Gallone, P., Romanazzi, G., Schena, L., Ippolito, A., and Salerno, M. G. 2005. Incidence of *Verticillium* wilt on olive in Apulia and genetic diversity of *Verticillium dahliae* isolates from infected trees. *J. Plant Pathol.* 87:13-23.
34. Peever, T. L., Canihos, Y., Olsen, L., Ibañez, A., Liu, Y. C., and Timmer, L. W. 1999. Population genetic structure and host specificity of *Alternaria* spp. causing brown spot of Minneola tangelo and Rough lemon in Florida. *Phytopathology* 89:851-860.
35. Pegg, G. F., and Brady, B. L. 2002. *Verticillium* Wilts. CAB International, Wallingford, UK.
36. Qin, Q. M., Vallad, G. E., Wu, B. M., and Subbarao, K. V. 2006. Phylogenetic analyses of phytopathogenic isolates of *Verticillium* spp. *Phytopathology* 96:582-592.
37. Raayaree, P., Ospina-Giraldo, M. D., Kang, S., Bath, R. G., Subbarao, K. V., Grant, S. J., and Dobinson, K. F. 2005. Mutations in *VMK1*, a mitogen-activated protein kinase gene, affect microscerotia formation and pathogenicity in *Verticillium dahliae*. *Curr. Genet.* 48:109-116.
38. Resende, M. L. V., Flood, J., and Cooper, R. M. 1994. Host specialization of *Verticillium dahliae*, with emphasis on isolates from cocoa (*Theobroma cacao*). *Plant Pathol.* 43:104-111.
39. Rowe, R. C. 1995. Recent progress in understanding relationships between *Verticillium* species and subspecific groups. *Phytoparasitica* 23:31-38.
40. Ryder, E. J., De Vos, N. E., and Bari, M. A. 1983. The Globe artichoke (*Cynara scolymus* L.). *HortScience* 18:646-653.
41. Schnathorst, W. C., and Mathre, D. E. 1966. Host range and differentiation of a severe form of *Verticillium albo-atrum* in cotton. *Phytopathology* 56:1155-1161.
42. Shah, D. A., and Madden, L. V. 2004. Nonparametric analysis of ordinal data in designed factorial experiments. *Phytopathology* 94:33-43.
43. Strausbaugh, C. A., Schroth, N. M., Weinhold, A. R., and Hancock, J. G. 1992. Assessment of vegetative compatibility of *Verticillium dahliae* tester strains and isolates from California potatoes. *Phytopathology* 82:61-68.
44. Subbarao, K. V., Chassot, A., Gordon, T. R., Hubbard, J. C., Bonello, P., Mullin, R., Okamoto, D., Davis, R. M., and Koike, S. T. 1995. Genetic-relationships and cross pathogenicities of *Verticillium dahliae* isolates from cauliflower and other crops. *Phytopathology* 85:1105-1112.
45. Tjamos, E. C. 1981. Virulence of *Verticillium dahliae* and *Verticillium albo-atrum* isolates in tomato seedlings in relation to their host of origin and the applied cropping system. *Phytopathology* 71:98-100.
46. Tsrar, L., Hazanovsky, M., Mordechi-Lebiush, S., and Sivan, S. 2001. Aggressiveness of *Verticillium dahliae* isolates from different vegetative compatibility groups to potato and tomato. *Plant Pathol.* 50:477-482.
47. Zeise, K., and von Tiedemann, A. 2002. Application of RAPD-PCR for virulence type analysis within *Verticillium dahliae* and *V. longisporum*. *J. Phytopathol.* 150:557-563.
48. Zeise, K., and von Tiedemann, A. 2002. Host specialization among vegetative compatibility groups of *Verticillium dahliae* in relation to *Verticillium longisporum*. *J. Phytopathol.* 150:112-119.